



RumiLick Bovine 50

Proteïn, spoormineraal aanvulling vir Beeste
Protein, trace mineral supplement for Cattle

Voedingstof	Eenheid / Unit		Nutrient
Ru-Proteïn (Min)	500	g/kg	Crude Protein (Min)
Ureum (Maks)	155	g/kg	Urea (Max)
Proteïn vanaf NPN (Maks)	89	%	Protein from NPN (Max)
Vog (Maks)	120	g/kg	Moisture (Max)
Kalsium (Min/Maks)	6 - 18	g/kg	Calcium (Min/Max)
Fosfaat (Min)	6	g/kg	Phosphorus (Min)
Magnesium (Min)	2	g/kg	Magnesium (Min)
Swael (Min)	8	g/kg	Sulphur (Min)
Sout (Min/Maks)	20 – 40%	g/kg	Salt (Min/Max)
Vitamiene A	16 000	IU/kg	Vitamin A
Mangaan	180	mg/kg	Manganese
Koper	65	mg/kg	Copper
Kobalt	1.2	mg/kg	Cobalt
Yster	250	mg/kg	Iron
Jodium	4	mg/kg	Iodine
Sink	250	mg/kg	Zinc
Selenium	0.8	mg/kg	Selenium
WAARSKUWINGS		WARNINGS	
Hierdie produk bevat NPN bronne - lees die NPN waarskuwings voordat die produk gebruik word		This product contain NPN sources – read the NPN warnings before using the product	



PRODUK BESKRYWING		PRODUCT DESCRIPTION	
<p>Bovine 50 is 'n proteïen, spoormineraal aanvulling vir beeste tydens die droë seisoen. Die gebruik van hierdie produk sal die diere help om massa en kondisie te handhaaf op weidings wat laag is in voedingswaarde en verteerbaarheid, wat 'n positiewe effek op reproduksie en winsgewendheid sal hê.</p>		<p>Bovine 50 is a protein, trace mineral supplement for cattle during the dry season. Using this product will assist animals to maintain their body weight and condition while on grazing that are deficient in nutritional value, and low in digestibility. This will increase their chances on conception, and increase herd profitability.</p>	
VOER INSTRUKSIES		FEEDING INSTRUCTIONS	
<p>Begin met aanvulling sodra die kondisie van weiding versleg, wat gewoonlik begin gebeur vandat die gras saad skiet. Voorsien die lek so gereeld as moontlik, verkieslik daagliks. Sodra aanbevole innames gereeld oorskry word, kan dit 'n aanduiding wees dat weiding onvoldoende is, en die diere na nuwe kampe verskuif moet word.</p>		<p>Start supplementation when the quality of grazing starts to decline, which normally happens as soon as the grass produces seed. Supply licks daily if possible. When the recommended intake is exceeded frequently, it could be an indication that the grazing is insufficient and animals need to be moved to fresh paddocks.</p>	
AANBEVOLE INNAMES:		RECOMMENDED INTAKE:	
Beeste:	350 – 450g/dier/dag)	Cattle:	350 – 450g/animal/day