



RumiLick

Pro 40

Proteïen, spoormineraal aanvulling vir Herkouers
Protein, trace mineral supplement for Ruminants

V 24943

Voedingstof	Eenheid / Unit		Nutrient
Ru-Proteïen (Min)	400	g/kg	Crude Protein (Min)
Ureum (Maks)	105	g/kg	Urea (Max)
Proteïen vanaf NPN (Maks)	75.3	%	Protein from NPN (Max)
Vog (Maks)	120	g/kg	Moisture (Max)
Kalsium(Maks)	30	g/kg	Calcium (Max)
*Fosfaat (Min)	7	g/kg	*Phosphorus (Min)
Magnesium (Min)	5	g/kg	Magnesium (Min)
Swael (Min)	6	g/kg	Sulphur (Min)
Vitamiene A	20 000	IU/kg	Vitamin A
Mangaan	200	mg/kg	Manganese
Koper	75	mg/kg	Copper
Kobalt	1	mg/kg	Cobalt
Yster	250	mg/kg	Iron
Jodium	5	mg/kg	Iodine
Sink	300	mg/kg	Zinc
Selenium	1	mg/kg	Selenium
*Ten spyte daarvan dat hierdie produk nie as 'n mineraal aanvulling geklasifiseer word nie, bevat dit lae vlakke van fosfaat, maar voldoen nie aan die minimum vereistes van 'n mineraal aanvulling volgens Wet 36 van 1947 nie. Raadpleeg u tegniese adviseur vir die redes hiervoor.		* Although this product is not registered as a mineral supplement, it does contain low levels of phosphorus, but does not comply to the minimum requirements of Act 36 of 1947 for a mineral supplement. Ask your technical advisor about the reasons for this.	
WAARSKUWINGS		WARNINGS	
Hierdie produk bevat NPN bronne - lees die NPN waarskuwings voordat die produk gebruik word		This product contain NPN sources – read the NPN warnings before using the product	



PRODUK BESKRYWING		PRODUCT DESCRIPTION	
Rumilick Pro 40 is 'n proteïen, spoormineraal aanvulling vir skape en beeste tydens die droë seisoen. Die aanvulling van diere met hierdie produk sal die diere help om massa en kondisie te handhaaf op weidings wat laag is in voedingswaarde en verteerbaarheid, wat 'n positiewe effek op reproduksie en winsgewendheid sa hê.		Rumilick Pro 40 is a protein, trace mineral supplement for sheep and cattle during the dry season. Supplementing animals with this product will assist them to maintain their body weight and condition while on grazing that are deficient in nutritional value, and low in digestibility. This will increase their chances on conception, and increase herd profitability.	
VOER INSTRUKSIES		FEEDING INSTRUCTIONS	
Begin met aanvulling sodra die kondisie van weiding versleg, wat gewoonlik begin gebeur vandat die gras saad skiet. Voorsien die lek so gereeld as moontlik, verkieslik daaglik. Sodra aanbevole innames gereeld oorskry word, kan dit 'n aanduiding wees dat weiding onvoldoende is, en die diere na nuwe kampe verskuif moet word		Start supplementation when the quality of grazing starts to decline, which normally happens as soon as the grass produces seed. Supply licks daily if possible. When the recommended intake is exceeded frequently, it could be an indication that the grazing is insufficient and animals need to be moved to fresh paddocks.	
AANBEVOLE INNAMES:		RECOMMENDED INTAKE:	
Skape:	80 – 100g/dier/dag	Sheep:	80 – 100g/animal/day
Beeste:	500 – 600g/dier/dag)	Cattle:	500 – 600g/animal/day