

NutriCam

Dry Sow  
& Boar



Klas: Volledige Droë Sog- & Beervoer / Class: Complete Dry Sow & Boar Feed

### V30180

Voedingstof	Eenheid / Unit		Nutrient
Proteïen (Min)	125	g/kg	Protein (Min)
Lisien (Min)	6.0	g/kg	Lysine (Min)
Metionien (Min)	2.0	g/kg	Methionine (Min)
Vog (Maks)	120	g/kg	Moisture (Max)
Vet (Min)	25	g/kg	Fat (Min)
Vesel (Maks)	80	g/kg	Fibre (Max)
Kalsium (Min/Maks)	67.5-12.0	g/kg	Calcium (Min/Max)
Fosfor (Min)	5.0	g/kg	Phosphorus (Min)
Sout (Maks)	5.0	g/kg	Salt (Max)

### PRODUKTESKRYWING / PRODUCT DESCRIPTION

'n Volledige voer vir dragtige soggies en sôe om goeie fetale ontwikkeling te verseker, terwyl die ideale liggaamskondisie terselfdertyd behou word. Dit kan ook as 'n volledige voer vir werkende bere gevoer word.

A complete feed for gestating gilts and sows to ensure good fetal development while at the same time maintaining the ideal body composition. It can also be fed as a complete feed for working boars.

### VOERINSTRUKSIES / FEEDING INSTRUCTIONS

Voer dragtige soggies en sôe na spening tot met jonging. Pas die voedingsvlakke aan om die verlangde liggaamskondisie te bereik, wat benodig word by jonging deur van die aanbevole inname-ryglyne gebruik te maak, soos hieronder aangedui. Die PorCare Pre-Farrow dieet moet vir die laaste 7 tot 10 dae voor jonging gevoer word.

Feed gestating gilts and sows from after weaning until farrowing. Adjust the feeding level to achieve the desired body condition required at farrowing using the recommended intake guidelines below. The PorCare Pre-Farrow diet should be fed for the last 7 to 10 days before farrowing.

NutriCam

Dry Sow  
& Boar



Klas: Volledige Volledige Droë Sog- & Beervoer / Class: Complete Dry Sow & Boar Feed

### AANBEVOLE INNAMES / RECOMMENDED INTAKE

Voer bere 2,0 - 2,5 kg per dag, afhangende van die grootte van die liggaam, kondisie en werkstempo. Voeders moet altyd skoon gehou word en vars water moet ten alle tye beskikbaar wees.	Feed boars 2,0 - 2,5 kg per day depending on body size, condition and work rate. Feeders must always be kept clean and water must always be available.		
Voer bere 2,0 - 2,5 kg per dag, afhangende van die grootte van die karkas, kondisie en werkstempo.	Feed boars 2,0 to 2,5 kg per day depending on body size, condition and work rate.		
	Gestation Period / Dragtigheidsperiode		
	0 - 28 days kg/d	28 - 90 days kg/d	90 days - farrow house kg/d
Gilts mated in optimum weigh range / Soggies gepaar in optimale gewigsklasse	1,6	1,8	2,7
Gilts heavier than 154 kg at mating / Soggies swaarder as 154 kg by paring	1,8	1,6	1,6
Thin sows / Maer sôe	2,7	2,7	2,7
Normal sows / Normale sôe	2,3	1,8	2,7
Fat sows / Vet sôe	1,6	1,6	1,6